

#### INSTRUCTIONS

PREPARATION: Shake well before use. Can be applied in combination with compatible plant growth regulators, pesticides, or other liquid fertilizers. If compatibility is in question, jar test a small quantity. Do not foliarly apply this product in concentrations greater than 10% without a preliminary foliar test.

APPLICATION: Designed for soil and/or foliar application. This product supplies nutrients and/or biostimulants to plants and should be applied when tissue and soil analysis identifies the need and proper rate of application for this product. Best results will be obtained when application is concentrated in the active root zone or on the plant surface. Apply foliar sprays with sufficient water to ensure uniform coverage without running off leaf surfaces. For maximum results, apply early in the morning or in the evening when air temperatures are optimum for foliar application. Full label rates can be applied as often as every 7–10 days, as needed.

**CONDITIONS:** Do not apply nutrients during the hottest time of the day or to plants under drought or flood stress. Application timing, intervals, and rates may vary according to individual plant requirement, stage of development, available nutrient levels in the soil, and overall nutritional status of the plants.

**CONSULT:** Consult your local Huma® Representative or agricultural/turf/horticultural specialist for use-specific recommendations.

STORAGE & DISPOSAL: Do not store this product below 50°F (10°C) or above 90°F (30°C). Keep product in original container. Do not transfer into food or drink containers. Triple rinse container when empty for recycling. Always dispose of container in accordance with local, state, and/or federal regulations.

**DISCLAIMER:** This product is intended as a supplement to a regular fertilization program and will not, by itself, provide all the nutrients normally required by plants.

# MAX PAK®

# **GUARANTEED ANALYSIS**

Sulfur (S)	3.00%
Boron (B)	0.60%
Cobalt (Co)	0.05%
Copper (Cu)	1.00%
Iron (Fe)	2.00%
Manganese (Mn)	1.00%
Molybdenum (Mo)	0.05%
Zinc (Zn)	3.50%
Dariyad from Baric Acid Cabal	t Sulfate Sadium Malyhdate

Derived from Boric Acid, Cobalt Sulfate, Sodium Molybdate, Copper Sulfate, Ferrous Sulfate, Manganese Sulfate, and Zinc Sulfate.

ALSO CONTAINS BENEFICIAL SUBSTANCES: 6.0% Organic Matter

### SHAKE WELL BEFORE USE

Do not foliarly apply this product in concentrations greater than 10% without a preliminary foliar test.

Full label rates can be applied as often as every 7 to 10 days, as needed.

WARNING: This product contains boron (B), which may be injurious to certain crops. The use of this fertilizing material on any crop(s) other than those recommended may result in serious injury to the crop(s). WARNING: Application of this material in excess may result in forage crops containing levels of molybdenum (Mo.) that are toxic to ruminant animals.

# **APPLICATION**

METHOD	SUGGESTED RATE		
	Field Crops	Tree or Vine Crops	Turf or Horticulture
Foliar band application at 50% overage	Up to 1 pt/ac, 1.25 L/ha	-	Up to 1 oz/1000 ft² 35 mL/100 m²
Foliar broadcast or sprinklers: solid, set, linear, or pivot (100% speed)	Up to 1 qt/ac,	Up to 2 qt/ac,	Up to 2 oz/1000 ft <sup>2</sup>
	2.5 L/ha	5 L/ha	70 mL/100 m <sup>2</sup>
Soil banded or injected through drip tape or micro sprinklers	Up to 1 qt/ac,	Up to 2 qt/ac,	Up to 2 oz/1000 ft <sup>2</sup>
	2.5 L/ha	5 L/ha	70 mL/100 m <sup>2</sup>
Soil broadcast spray incorporated, flood or furrow irrigated	Up to 2 qt/ac,	Up to 1 gal/ac,	Up to 3 oz/1000 ft <sup>2</sup>
	5 L/ha	10 L/ha	105 mL/100 m <sup>2</sup>

10.85 lb per gallon @ 68°F. 1.30 kg per liter @ 20°C. pH 1.0-2.0 2.5 gallons • Net Wt. 27.13 lb 9.46 liters • Net Wt. 12.30 kg Lot No. 12345678

huma.us

#### SAFETY

#### KEEP OUT OF REACH OF CHILDREN

Information regarding the contents and levels of metals in this product is available on the Internet. http://www.aapfco.org/metals.html

The liquid and mists may cause irritation to the eyes, skin, and respiratory tract. This product may be toxic by ingestion or inhalation of high mist concentrations. The NIOSH IDLH for manganese compounds is 500mg/m³ (as Mn).

CLASSIFICATION: HAZARD CATEGORY 2 
SIGNAL WORD: N/A

**HAZARD STATEMENT:** H411 Toxic to aquatic life with long lasting effects.

PRECAUTIONARY STATEMENT: P273 Avoid release to the environment if this is not the intended use; P391 Collect spillage; P501 Dispose of contents/containers in accordance with federal, state, and local laws regulating fertilizers.

PRODUCT CHARACTERISTICS: A clear to slightly hazy, dark green, liquid having a sweet citrus-type odor. The color, pH, specific gravity, as well as the smell, can vary as this product is a formulation made with organic and organically active ingredients. Variations may occur due to changes in temperature, atmospheric pressure, weather conditions, and/or exposure to light from when the product leaves the factory until it reaches the final consumer.

CONDITIONS OF SALE: The information contained on this label is believed to be accurate and reliable. Buyer and user acknowledge and assume all liability resulting from the use of this material. Follow directions carefully. Timing, method of application, weather, plant and soil conditions, and other factors are beyond the control of the seller.

This product is powered by Micro Carbon Technology®.

© 2023 by Huma, Inc. 1331 W. Houston Ave., Gilbert, AZ 85233 480.961.1220 Publication No. HP&S-292-230921